

ORIGINAL ARTICLE

DOI: XXX

Effects of altitude on biceps brachii and erector spinae muscles oxygen saturation during basic cardiopulmonary resuscitation: a simulation study

Sonia Piñero-Saez¹, Angel López-González^{1*}, Isabel María Guisado-Requena¹, Jaime López-Tendero², Carmen María Guerrero-Agenjo², Francisco García-Alcaraz¹, Ignacio Martínez-González-Moro³, Joseba Rabanales-Sotos¹

1. Nursing Faculty of the Albacete, Castilla-La Mancha University, Albacete, Spain.

2. Castilla-La Mancha Health Service, Albacete, Spain.

3. Institute for Researching in Aging, Murcia, Spain.

*Corresponding author: Angel Lopez Gonzalez; Email: angel.lopez@uclm.es

Published online: 2024-09-15

Abstract: **Objective:** To assess biceps brachii and erector spinae muscular oxygen saturation (SmO₂) by near infrared spectroscopy (NIRS), during 10 minutes of resuscitation at simulated altitudes of 600, 3000 and 5000 m before and after carrying out a simulation program for adaptation to hypoxia. Performing and maintaining a good-quality cardiopulmonary resuscitation (CPR) at higher altitudes may pose a significant challenge to resuscitators due to decrease in arterial oxygen saturation. This fact adversely effects the quality of resuscitation.

Methods: Participants performed 10 minutes of CPR on a mannequin in the laboratory in environments that simulated altitudes. Subsequently, a standardized altitude conditioning protocol was carried out using intermittent hypoxia. The participants performed CPR again under the conditions and altitudes previously referred to.

Results: Initial heart rate (HR) at 5000 > 3000 m, and both > 600 m. HR at each altitude was higher conditioning at the end of CPR. The SmO₂ of both muscles showed no differences at the beginning and at the end of CPR and was higher in both muscles after the conditioning program before and at the end of CPR. In both muscles, SmO₂ values before and after conditioning show a slightly increasing trend during CPR.

Conclusion: NIRS use allows developing an optimum training plan. The rescuer will know his limits and optimize his performance. The improvement in physical performance and recovery capacity induced by intermittent hypoxia conditioning programs increases the quality of CPR in prolonged cardiac arrests and in adverse conditions, such as at high altitudes.

Keywords: Basic Cardiac Life Support; Heart Massage; High Altitude; Hypoxia; Near-Infrared Spectroscopy

Cite this article as: Piñero-Saez S, López-González A, Guisado-Requena IM, et al. Effects of altitude on biceps brachii and erector spinae muscles oxygen saturation during basic cardiopulmonary resuscitation: a simulation study. *Front Emerg Med*. In Press.

1. Introduction

Survival rates of out of hospital cardiac arrests (OHCA) is estimated to be below 10% (1). Good-quality cardiopulmonary resuscitation (CPR) maneuvers benefit survival after OHCA —by even doubling or tripling the survival rate (1,2). During CPR, chest compressions (CC) combined with breaths following strategy 30:2 are potentially effective in helping survival of people with OHCA (3). For this purpose, adult patients require CC between 5 and 6 centimeters (cm) deep, in 50:50-proportion compression-decompression cycles achieved through compressions exerted mid-sternum at a rate of 100 to 120 per minute (4). During CPR maneuvers, biceps brachii and erector spinae muscles play a significant role in the position and performance of CC (5).

It has been proven that in the general population and under ideal environmental conditions, the resuscitator's efforts

during CPR are essentially aerobic and submaximal, and the cardiorespiratory system can tolerate these exercises well. The resuscitator's shape influences CPR quality as well as endurance until the arrival of help (6,7).

In recent years people have shown a considerably higher interest in outdoor activities at a high altitude, such as hiking, mountain climbing and skiing. Even the number of amateurs becoming fans of such disciplines has grown noticeably in the past years (8) —it is common to find sport teams exercising or supporters following sport events over medium and high altitudes. Staying and moving at a high altitude produces a variety of physiological effects, such as an increase in cardiac and respiratory workload, mainly during physical activity (9,10).

Performing and maintaining a good-quality CPR at higher altitudes may pose a significant challenge to resuscitators. Pre-

vious studies have demonstrated that after a 5-minute CPR at altitude, resuscitators' arterial oxygen saturation (SpO₂) decreases and heart rate (HR) increases. This fact adversely influences CC and ultimately, CPR quality (11–16).

Endurance of the skeletal muscle relies on oxidative metabolism and thus on the amount of oxygen (O₂) available. The percentage of O₂ absorbed by muscular tissue during oxidative metabolism for generating power can be calculated through muscular oxygen saturation (SmO₂) (17,18).

It is possible to estimate tissue oxygenation by near infrared spectroscopy (NIRS) (18). NIRS establishes SmO₂ based on the relation between oxyhemoglobin and hemoglobin total values in blood and expresses it as a percentage (%SmO₂) (19). NIRS is considered an excellent technique to evaluate performance during physical activity given the proportional relationship between %SmO₂ and the ability to carry out physical activity (20,21).

NIRS has proved to show fatigue better than HR-based measurements (13). It has been demonstrated as well that correlation between NIRS and maximal lactate steady state is a valid sign of endurance and training (22,23).

NIRS devices connected to displays (computers, tablets, or watches) allow real-time measuring of SmO₂ consumed by the body during physical activity, outdoors and in laboratory tests (24,25). In sports, this muscular-oxygenation-based measuring method has been employed as a predictor of endurance in disciplines such as climbing, race walking and cycling (26,27).

Normobaric intermittent hypoxia exposure (NIHE) occurs when breathing air with low O₂ content under regular environmental pressure conditions, with the aim to stimulate hematological adjustments (increased production of erythropoietin, hemoglobin, erythrocytes) and changes in performance (modifications in pH regulation and lactate transport) which are typical of acclimation to altitude (28).

No previous studies have monitored the SmO₂ of biceps brachii and erector spinae during CPR at different altitudes or studied the likely effect of a program for adaptation to hypoxia caused by altitude. Thus, our objectives were: a) to assess biceps brachii and erector spinae SmO₂ during 10 minutes of CC at simulated altitudes of 600, 3000 and 5000 meters (m) after running a program for adaptation to hypoxia caused by altitude; b) to describe the effects of altitude training on biceps brachii and erector spinae SmO₂ related to CPR at such altitudes.

2. Methods

2.1. Study design

An analytical before-after study in which participants, selected for convenience performed CC inside a laboratory for 10 minutes on a mannequin in environments simulating different altitudes (600, 3000 and 5000 m). It was hence followed by an intervention based on a standardized protocol for adaptation to altitude by inducing intermittent hy-

poxia. After acclimation to hypoxia, the same participants performed CC one more time under the same simulated conditions and altitudes.

2.2. Study population

10 people in their third or fourth year of bachelor's degree in nursing (5 women) with training in CPR in compliance with resuscitation guidelines ERC 2021 (ERCGR2021) were invited to participate (4) —all of them being of age and none suffering from any illness or physical disability preventing them to participate in the study.

Exclusion criteria were suffering from any chronic or acute condition, serious psychiatric disorder, tumor pathology, uncontrolled hypertension, serious imbalance problems, intolerance to hypoxia caused by the different simulated altitudes during CPR, inability to reach 85% of correction in CC in CPR at 600 m. Three participants were excluded as they did not comply with the last criterion.

2.3. Study process

Sociodemographic variables (sex, age, and level of education) as well as the weight and height of each participant was collected initially. All of them took the primary test on tolerance to hypoxia according to protocol iAltitude (29). The flowchart of the study is shown in figure 1.

- Test 1: All participants performed 10 minutes of uninterrupted CPR on the mannequin placed on the floor (CC to ventilation by bag-valve mask ratio=30:2) at simulated altitudes of 600, 3000 and 5000 m. Prior to CPR at each altitude, participants had a 10-minute period for adaptation to fraction of inspired oxygen (FiO₂) as it was expected to be in each simulated environment. In between tests 60 minutes served for physical recovery.

- NIHE physical conditioning: Following the initial measurements, an intervention was carried out alternating phases of hypoxia and normoxia according to the iAltitude conditioning program, with 967 minutes distributed in sessions of 56–72 minutes, alternating cycles of hypoxia (FiO₂ of 0.14 in the initial sessions to 0.11 in the final ones) and normoxia, resulting in 728 minutes of hypoxia total time.

- Test 2: Once the conditioning program for intermittent hypoxia was completed, all participants performed 10-minute continued CPR at simulated altitudes of 600, 3000 and 5000 m with a previous 10-minutes period for adaptation to FiO₂ in accordance with the simulated altitude, and 60 minutes between tests for physical recovery.

Considering the advice by Bastida-Castillo *et al.* (30) about where to place the device to detect changes in oxygenation in muscular tissue, each participant had two Humon Hex® devices placed on their dominant body zones. One device for the biceps brachii, on the wider area of the arm, and the other for the erector spinae, placed at the center of the muscle. Each Humon Hex® was synchronized with an iPad®, Apple Inc. by switching on the corresponding application to check SmO₂, amount of time of exercise and HR (25). SmO₂




Figure 1 Study flow diagram

Table 1 Basic characteristics of the population

	Gender (Men 4, women 3)	Mean	SD	CV	t-value	P-value
Age, years	Men	23.3	4.4	18.26		
	Women	22.4	1.6	7.24	1.356	0.096
Weight, Kg	Men	80.64	11.36	14.09		
	Women	64.86	9.35	14.42	3.393	0.002
Height, cm	Men	174.2	7.63	4.35		
	Women	163.6	7.63	4.61	2.810	0.006
BMI, Kg/m ²	Men	21.1	2.66	10.16		
	Women	18.1	2.55	10.82	2.243	<0.001
HR, bpm	Men	82.8	14.4	18.02		
	Women	80	10.5	13.36	0.230	0.410
SBP, mmHg	Men	135.5	14.1	10.63		
	Women	123.0	15.3	12.44	1.875	0.390
DBP, mmHg	Men	77.7	10.0	12.87		
	Women	79.3	9.3	11.73	0.369	0.358
SpO ₂ , %	Men	99.3	0.4	0.4		
	Women	99.0	0.8	0.81	0.000	0.500

BMI: Body mass index; HR: Heart rate; SBP: Systolic pressure; DBP: Diastolic pressure; SpO₂: Partial oxygen saturation; bpm: Beats per minute; cm: Centimeter; kg: Kilogram; SD: Standard deviation; CV: Coefficient of variation
In bold type: $P \leq 0.05$

values were concluded from the SmO₂/exercise time curves generated by Humon Hex®, from which data were also selected for subsequent phases: a) prior to exercise; b) drop phase: when SmO₂ drops drastically (Humon Hex® indicates this fact with a change of color (orange); c) maximum

drop coinciding with exercise maximum intensity and maximum peak or maximum oxygen uptake, (red color); and d) recovery phase: after the exercise, when values return to normal (indicated by another color change color on the graph from blue to green).



Figure 2 %SmO₂ evolution during CPR; a) biceps brachii; b) erector spinae and quality of CC at different altitudes before and after conditioning

The CPR test was performed on an electronic mannequin designed for CPR education purposes (Laerdal Resusci Anne

CPR-D and SkillReporter®; Medical Laerdal; Stavanger, Norway) placed on the ground and configured according to the

Table 2 Comparison of mean values of HR, SmO2 (%) in biceps brachii and erector spinae during CPR at different altitudes before and after conditioning

			F-value	P-value
Pre-conditioning values	HR, bpm	Altitude	3.144	0.080
		Moments	23.198	0.003
	Biceps brachii SmO2, %	Altitude	0.268	0.769
		Moments	3.448	0.113
	Erector spinae SmO2, %	Altitude	2,138	0,180
		Moments	1.797	0.251

HR: Heart rate; SmO2: Muscle oxygen saturation; Moments: CPR initial, CPR final;
bpm: Beats per minute
In bold type $P \leq 0.05$ statistically significant

Table 3 Comparison of HR and SmO2 mean values (%) in biceps brachii and erector spinae during CPR at 600 m with those obtained at 3000 and 5000 m (preconditioning)

600 m variable	Moment	Mean value	Altitude	Moment	Mean value	t-value	P-value
HR, bpm	Initial	81.60	3000	Initial	90.5	0.579	0.584
	Final	169.1		Final	153.7	1.778	0.126
			5000	Initial	92.7	2.379	0.055
				Final	163.35	5.712	0.001
Biceps brachii SmO2, %	Initial	56.85	3000	Initial	54.59	0.452	0.667
	Final	58.28		Final	60.94	0.732	0.492
			5000	Initial	61.14	0.667	0.530
				Final	60.75	0.430	0.682
				Final	65.12	2.047	0.087
Erector spinae SmO2, %	Initial	72.79	3000	Initial	59.96	2.333	0.080
	Final	74.47	5000	Final	62.30	1.854	0.113
				Initial	66.34	1.675	0.169
				Final	65.12	2.047	0.087

HR: Heart rate; SmO2: Muscle oxygen saturation; bpm: Beats per minute
In bold type $P \leq 0.05$ statistically significant

Table 4 Comparison of biceps brachii and erector spinae SmO2 at the time of CPR at different altitudes before and after conditioning

		F-value	P-value
Biceps brachii SmO2, %	Altitude	0.384	0.689
	Moment	12.644	0.012
	Phase	0.236	0.644
	Altitude-moment	1.204	0.334
	Altitude-phase	2.570	0.118
	Phase-moment	4.921	0.068
	Altitude-moment-phase	2.427	0.130
Erector spinae SmO2, %	Altitude	0.311	0.744
	Moment	20.777	0.020
	Phase	10.261	0.049
	Altitude-moment	15.349	0.004
	Altitude-phase	6.250	0.034
	Phase-moment	50.314	0.006
	Altitude-moment-phase	9.992	0.012

SmO2: Muscle oxygen saturation; Moment: initial and final CPR; Phase: pre and post conditioning
In bold type $P \leq 0.05$ statistically significant

ERC Guidelines 2021 (4). The strength necessary to perform thorax compressions with a depth of 5 cm in this device was >28.5 kp. CC quality was obtained from the manikin software and is the result of integrating CC with adequate chest depth, rhythm, and relaxation.

iAltitude Simulator® (Madrid, España) allowed both the simulation of different altitudes (600, 3000 and 5000 m) and the physical conditioning program. It includes a circuit for hypoxia induction and an adjustable face mask (individual use). The FiO2 was 0.21, 0.14 and 0.11 for 600, 3000 and 5000 m, re-

spectively.

Resuscitators' physiological demands during tests were registered using the software iAltitude Trainer® (Madrid, España). SpO₂ was measured with a pulse oximeter (Oximeter DF8 W / USB 3018L XPOD, ear clip SNRS 8000Q2; Nonin Medical, Plymouth, EE. UU.) placed on the right earlobe of each participant. HR was measured with a sensor Polar Team H7 (Kempele, Finland) located above the xiphoid process of the sternum (31). A blood pressure monitor was placed on the right arm to monitor BP (OMRON M5- I, Matsusaka, Japan). During the resting phase, each participant had an electrocardiogram, and HR and blood pressure were tested.

2.4. Statistical analysis

Quantitative variables have been described using mean values, standard deviation (SD) and coefficient of variation (CV); absolute frequency and percentages have been used for qualitative variables. The Shapiro–Wilk and Levene tests have been used to check the normal distribution of data and to identify homogeneity of variances. The comparison of means of dependent variables was performed using the paired t-test and independent variables with Student's t test. Practical significance was assessed by calculating the size of Cohen's d-effect for dependent variables between baseline and end-to-end data.

Effect sizes (d) greater than 0.8, between 0.8 and 0.5, between 0.5 and 0.2, and less than 0.2 were considered large, moderate, small, and trivial, respectively (32). Using the analysis of variance of repeated samples (ANOVA), the differences in the values of HR, (%) SmO₂ in biceps brachii and erector spinae at the different altitudes before and after conditioning were compared. A statistical relevance was considered only when $P \text{ value} \leq 0.05$. All the statistical analyses were performed with IBM® SPSS® Statistics v28.

2.5. Ethical considerations

All participants approved to participate with written informed consent. This study respected the ethical principles of the Declaration of Helsinki. The protocol for the study was approved by the Ethics Committee for Investigation with medicinal products of the Integrated Care Management of Albacete, Spain (code 2021-094).

3. Results

Table 1 shows the basic characteristics of the participants comparing both genders. Usual anthropometric differences between both genders are observed with higher values in the weight, height, and Body mass index (BMI).

Neither of the two muscles showed significant differences in SmO₂ at the end of CPR performed at different altitudes before and after conditioning. On the other hand, HR during CPR did show significant differences between the initial and final values after the intervention, but not between altitudes (Table 2).

Table 3 shows the comparison of HR and SmO₂ values in bi-

ceps brachii and erector spinae during CPR at 600 m with the values obtained at 3000 and 5000 m prior to the conditioning program. There were no differences in SmO₂ values of either of the muscles when compared with values at 600 m. Relevant differences ($P < 0.001$) were seen only in HR after CPR at 5000 m (169.1 vs 163.35 at 600 m).

On the other hand, when analyzing the variance of repeated samples considering the effect of conditioning (Table 4), it is obtained that the significant differences are associated in the biceps brachii with the moment in which the SmO₂ is obtained (beginning or end of the exposure). In the erector spinae, differences are also observed between before and after conditioning and when the three situations are combined.

Figure 2 shows evolution of biceps brachii and erector spinae SmO₂ and quality CC during CPR at different altitudes before and after conditioning.

It is noticeable that, in both muscles, SmO₂ values before and after conditioning show a slightly rising trend during CPR, and %SmO₂ values are lower at 5000 m. In pre-conditioning, they were 82% at 600 and 5000 m and 89% at 3000. In post-conditioning, the results were 86%, 87% and 89% at 600, 3000 and 5000 m respectively.

4. Discussion

This study analyzed the SmO₂ in biceps brachii and erector spinae muscles during a 10-minute CPR at simulated altitudes of 600, 3000 and 5000 m, and the influence of an intervention based on a training program on the SmO₂ of muscles and on said altitudes during CPR maneuvers (with help from device Humon Hex®). CPR has been performed on a standardized mannequin with invariable resistance for both muscles independent of the altitude at which CPR was performed (6). The results of this study can be used to better understand the mechanism of how each muscle works during CC and why fatigue sets in.

During preconditioning, HR was higher after CPR at all altitudes, which is reasonable after every effort in physical activity. SmO₂ in both muscles followed a similar pattern: it was lower at all altitudes at the beginning of CPR. SmO₂ at 600 and 3000 m increased (33–34), however this increase did not recur after CPR at 5000 m, which could be due to a decrease in CC quality, by compressions not being deep enough and thus lower the efforts of the resuscitator, as reported by McDonnals et al. (35).

Over increasing physical effort, SmO₂ decreases as reported by Wilkinson et al. (36). In our study, biceps brachii and erector spinae SmO₂ did not decrease over CPR performance: effort intensity remains constant while performing CC in compliance with quality criteria; increasing intensity would reduce quality, as it was stated before.

The absence of significant differences when comparing SmO₂ at 600 m before and after CPR to SmO₂ measurements at 3000 and 5000 m could be reasonable as CPR is a medium effort activity for a medium sized individual and because

CC quality depends greatly on BMI amongst all other factors (7,37–39).

Following the conditioning program, before and after CPR, SmO₂ exceeded the values obtained before conditioning in both muscles. Likewise, in our study, following the conditioning program, except for 5000 m altitude, SmO₂ in both muscles at 600 and 3000 m was higher than at the end of CPR than at the beginning. This information matches the reports by Wilkinson et al. about physiological demands meaning an increasing effort (35). In our study, the increase in demands were caused by altitude hypoxia, not because of CC efforts. Therefore, we can declare that a conditioning program for hypoxia improves tissue oxygenation in scenarios of physical activity efforts in altitudes (10,40).

Finally, as already stated by Jui-Yi T et al. and proven in our study too, when comparing biceps brachii and erector spinae SmO₂ at 10-minute CPR, before and after conditioning, we notice similar SmO₂ values, as both muscles participate equally in the execution of CC (5).

The outcomes from this study can be useful to better understand the participation of muscle groups in CC during CPR, and fatigue, which reduces the quality of CC. NIRS devices connected to screens (watch, tablet, etc.) may be applied to report resuscitators in real time about low oxygenation levels in muscles and when resuscitators need to switch places in order to maintain the quality of CC.

5. Limitations

As for limitations in our study, we would mention the small sample size, which was due to the complexity of the intervention and duration of the conditioning program for hypoxia. Moreover, unlike the biceps brachii muscles, the erector spinae muscle movement in CC might have interfered with registration and collection of SmO₂ values. Another limitation was lacking measurement of adipose tissue in the areas where portable NIRS devices were placed, since a higher amount of adipose tissue affects penetration of the source light and thus data analysis by the devices using NIRS as well.

6. Conclusion

We conclude that NIRS use allows developing an optimum training plan, making the resuscitator aware of the effort they can make without exceeding any limit and by optimizing performance.

The improvement in physical performance and recovery capacity induced by intermittent hypoxia conditioning programs increases the quality of CPR in prolonged cardiac arrests and in adverse conditions, such as at high altitudes.

7. Declarations

7.1. Acknowledgement

None.

7.2. Authors' contribution

SP-G: Conceptualization (equal), formal analysis (equal), methodology (equal), project administration (lead), software (equal), supervision (equal), writing review editing (lead); AL-G: Conceptualization (equal), methodology (equal), resources (equal); CMG-A: formal analysis (equal), methodology (equal), software (equal), supervision (equal), validation (equal); IMG-R: Conceptualization (equal), methodology (equal), resources (equal); CMG-A: formal analysis (equal), methodology (equal), software (equal), supervision (equal), validation (equal); JL-T: Conceptualization (equal), methodology (equal), project administration (lead), resources (equal), supervision (lead), writing review editing (lead). FG-A: Conceptualization (equal), data curation (lead), formal analysis (equal), investigation (equal), resources (equal), writing original draft (lead); IM-GM: Conceptualization (equal), data curation (lead), formal analysis (equal), investigation (equal), resources (equal), writing original draft (lead); JR-S: Conceptualization (equal), data curation (lead), formal analysis (equal), investigation (equal), resources (equal), writing original draft (lead).

7.3. Conflict of interest

The authors of this paper declare that there were no conflicts of interest.

7.4. Funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

References

1. Kim SY, Lee SY, Kim TH, Shin SD, Song KJ, et al. Location of out-of-hospital cardiac arrest and the awareness time interval : a nationwide observational study. *Emerg Med J*. 2022;39:118–23.
2. Semeraro F, Greif R, Böttiger BW, Burkart R, Cimpoesu D, Georgiou M, et al. European resuscitation council guidelines 2021: systems saving lives. *Resuscitation*. 2021;161:80–97.
3. Cunningham LM, Mattu A, O'Connor RE, Brady WJ. Cardiopulmonary resuscitation for cardiac arrest: the importance of uninterrupted chest compressions in cardiac arrest resuscitation. *Am J Emerg Med*. 2012;30(8):1630–8.
4. Olasveengen T, Semeraro F, Ristagno G, Castren M, Handley AJ, Kuzovlev A, et al. European resuscitation council guidelines for resuscitation 2021: basic life support. *Resuscitation*. 2021;161:98–114.
5. Jui-Yi T, Fong-Chin S, Pai-Chin T, Ming-Yuan H, Su-Chun C, Hsun-Wen C, et al. Electromyography activity of selected trunk muscles during cardiopulmonary resuscitation. *Am J Emerg Med*. 2014;32(3):216–20.
6. Fernandez-Lopez M, Mateos-Lorenzo J, Pavon-Prieto P, Freire-Tellado M, Sanchez-Santos L, Estany-Gestal A, et al. Chest compressions and cardiopulmonary resuscita-

- tion. Why do we need technical or physical training? Resuscitation. 2015;96(1):13.
7. López-González A, Sánchez-López M, Rovira-Gil E, Ferrer-López V, Martínez-Vizcaíno V. Influencia del índice de masa corporal y la forma física de jóvenes universitarios en la capacidad de realizar compresiones torácicas externas de calidad sobre maniquí. *Emergencias*. 2014;26:195-201. [Spanish].
 8. De Aquino-Lemos V, Antunes HK, Dos Santos RV, Lira FS, Tufik S, De Mello MT. High altitude exposure impairs sleep patterns, mood, and cognitive functions. *Psychophysiology*. 2012;49(9):1298-306.
 9. Savourey G, Launay JC, Besnard Y, Guinet A, Travers S. Normo- and hypobaric hypoxia: are there any physiological differences? *Eur J Appl Physiol*. 2003;89:122-6.
 10. Coppel J, Hennis P, Gilbert-Kawai E, Grocott MPW. The physiological effects of hypobaric hypoxia versus normobaric hypoxia: a systematic review of crossover trials. *Extrem Physiol Med*. 2015;4:2.
 11. Suto T, Saito S. Considerations for resuscitation at high altitude in elderly and untrained populations and rescuers. *Am J Emerg Med*. 2014;32:270-6.
 12. Wang JC, Tsai SH, Chen YL, Hsu CW, Lai KC, Liao WI, et al. The physiological effects and quality of chest compressions during CPR at sea level and high altitude. *Am J Emerg Med*. 2014;32:1183-8.
 13. Narahara H, Kimura M, Suto T, Saito H, Tobe M, Aso C, et al. Effects of cardiopulmonary resuscitation at high altitudes on the physical condition of untrained and unacclimatized rescuers. *Wilderness Environ Med*. 2012;23:161-4.
 14. Sato T, Takazawa T, Inoue M, Tada Y, Suto T, Tobe M, et al. Cardiorespiratory dynamics of rescuers during cardiopulmonary resuscitation in a hypoxic environment. *Am J Emerg Med*. 2018;36:1561-4.
 15. Carballo-Fazanes A, Barcala-Furelos R, Eiroa-Bermúdez J, Fernández-Méndez M, Abelaíras-Gómez C, Martínez-Isasi S, et al. Physiological demands of quality cardiopulmonary resuscitation performed at simulated 3250 meters high. *Am J Emerg Med*. 2020;38:2580-5.
 16. Egger A, Niederer M, Tscherny K, Burger J, Fuhrmann V, Kienbacher C, et al. Influence of physical strain at high altitude on the quality of cardiopulmonary resuscitation. *Scan J Trauma Resusc Emerg Med*. 2020;28(1):19.
 17. Inglis EC, Iannetta D, Murias JM. Evaluating the NIRS-derived microvascular O₂ extraction “reserve” in groups varying in sex and training status using leg blood flow occlusions. *Public Library of Science*. 2019;14(7):1-33.
 18. Ferrari M, Mottola L, Quaresima V. Principles, techniques and limitations of near infrared spectroscopy. *J Appl Physiol*. 2004;29(4):463-87.
 19. Özyener F. Evaluation of intra-muscular oxygenation during exercise in humans. *J Sports Sci Med*. 2002;1(1):15-9.
 20. Hamaoka T, McCully KK, Quaresima V, Yamamoto K, Chance B. Near-infrared spectroscopy/imaging for monitoring muscle oxygenation and oxidative metabolism in healthy and diseased humans. *J Biomed Opt*. 2007;12(6):1-16.
 21. Wilkinson TJ, White A, Nixon DG, Gould DW, Watson EL, Smith A. Characterizing skeletal muscle haemoglobin saturation during exercise using near-infrared spectroscopy in chronic kidney disease. *Clin Exp Nephrol*. 2019;23(1):32-42.
 22. Farzam P, Starkweather Z, Franceschini MA. Validation of a novel wearable, wireless technology to estimate oxygen levels and lactate threshold power in the exercising muscle. *Physiol Rep*. 2018;6(7):e13664.
 23. Bellotti C, Calabria E, Capelli C, Pogliaghi S. Determination of maximal lactate steady state in healthy adults: can NIRS help? *Med Sci Sports Exerc*. 2013;45(6):1208-16.
 24. Endo T, Kime R, Watanabe T, Fuse S, Murase N, Kurosawa Y, et al. Reduced optical path length in the vastus lateralis during ramp cycling exercise. *Adv Exp Med Biol*. 2020;1232(1):239-44.
 25. Humon. Humon Hex. 2023. Available at: <https://humon.es/> [Access 4 October 2023].
 26. Fryer SM, Stoner L, Stone KJ, Giles D, Sveen J, Garrido I, et al. Forearm muscle oxidative capacity index predicts sport rock-climbing performance. *Eur J Appl Physiol*. 2016;116(8):1479-84.
 27. Paredes-Ruiz MJ, Jódar-Reverte M, Ferrer-López V, Martínez-González-Moro I. Muscle oxygenation of the quadriceps and gastrocnemius during maximal aerobic effort. *Revista Brasileira de Medicina do Esporte*. 2021;27:212-27.
 28. Levine BD. Intermittent Hypoxic Training: fact and fancy. *High Alt Med Biol*. 2002;3(2):177-93.
 29. iAltitude. Protocolo precondicionamiento a hipoxia normobárica. Madrid: iAltitude; 2019. [Spanish].
 30. Bastida-Castillo A, Gómez-Carmona CD, Pino-Ortega J. Efectos del tipo de recuperación sobre la oxigenación muscular durante el ejercicio de sentadilla. *Kronos*. 2016;15(2):1-12. [Spanish].
 31. Hernández-Vicente A, Hernando D, Marín-Puyalto J, Vicente-Rodríguez G, Garatachea N, Pueyo E, Bailón R. Validity of the polar H7 heart rate sensor for heart rate variability analysis during exercise in different age, body composition and fitness level groups. *Sensors (Basel)*. 2021;21(3):902-16.
 32. Cohen J (1988). *Statistical power analysis for the behavioral sciences* 2nd ed. Hillsdale, NJ: Lawrence Erlbaum Associates, Publishers; 1988.
 33. Rosales MA, Shute RJ, Hailes WS, Collins CW, Ruby BC, Slivka DR. Independent effects of acute normobaric hypoxia and hypobaric hypoxia on human physiology. *Nature portfolio*. 2022;12:19570.
 34. Hansen J, Sander M, Hald CE, Victor RG, Thomas GD. Metabolic modulation of sympathetic vasoconstriction in human skeletal muscle: role of tissue hypoxia. *J Physiol*. 2000;527(2):387-36.

35. McDonald CH, Heggie J, Jones CM, Thorne CJ, Hulme J. Rescuer fatigue under the 2010 ERC guidelines, and its effect on cardiopulmonary resuscitation (CPR) performance. *Emerg Med J.* 2013;30:623-7.
36. Wilkinson TJ, White AEM, Nixon DGD, Gould DW, Watson EL, Smith AC. Characterising skeletal muscle hemoglobin saturation during exercise using near-infrared spectroscopy in chronic kidney disease. *Clin Exp Nephrol.* 2019;23:32-42.
37. Abelairas C, Romo V, Barcala R, Palacios A. Efecto de la fatiga física del socorrista en los primeros cuatro minutos de la reanimación cardiopulmonar posrescate acuático. *Emergencias.* 2013;25:184-90. [Spanish].
38. Hong DY, Park SO, Lee KR, Baek KJ, Shin DH. A different rescuer changing strategy between 30:2 cardiopulmonary resuscitation and hands-only cardiopulmonary resuscitation that considers rescuer factors: a randomized cross-over simulation study with a time-dependent analysis. *Resuscitation.* 2012;83(3):353-9.
39. Russo SG, Neumann P, Reinhardt S, Timmermann A, Niklas A, Quintel M, et al. Impact of physical fitness and biometric data on the quality of external chest compression: a randomized, crossover trial. *BMC Emerg Med.* 2011;11:20.
40. Sharma AP. Factors affecting sea-level performance following altitude training in elite athletes. *J Sci Sport Exerc.* 2022;4(4):315-30.